

Chemical Evolution of the Magellanic Clouds: analytical models

B.E.J. Pagel¹ and G. Tautvaišienė²

¹*NORDITA, Blegdamsvej 17, Dk-2100 Copenhagen Ø, Denmark (e-mail: pagel@nordita.dk)*

²*Institute of Theoretical Physics and Astronomy, Goštauto 12, Vilnius 2600, Lithuania (e-mail: taut@itpa.lt)*

Received; in original form.....

ABSTRACT

We have extended our analytical chemical evolution modelling ideas for the Galaxy (Pagel & Tautvaišienė 1995, 1997) to the Magellanic Clouds. Unlike previous authors (Russell & Dopita 1992; Tsujimoto *et al.* 1995; Pilyugin 1996), we assume neither a steepened IMF nor selective galactic winds, since among the α -particle elements only oxygen shows a large deficit relative to iron and a similar deficit is also found in Galactic supergiants. Thus we assume yields and time delays identical to those that we previously assumed for the solar neighbourhood. We include inflow and non-selective galactic winds and consider both smooth and bursting star formation rates, the latter giving a better fit to the age-metallicity relations. We predict essentially solar abundance ratios for primary elements and these seem to fit most of the data within their substantial scatter. Our LMC model also gives a remarkably good fit to the anomalous Galactic halo stars discovered by Nissen & Schuster (1997).

Our models predict current ratios of SNIa to core-collapse supernova rates enhanced by 50 per cent and 25 per cent respectively relative to the solar neighbourhood, in fair agreement with ratios found by Cappellaro *et al.* (1993) for Sdm-Im relative to Sbc galaxies, but these ratios are sensitive to detailed assumptions about the bursts and a still higher enhancement in the LMC has been deduced from X-ray studies of remnants by Hughes *et al.* (1995). The corresponding ratios integrated over time up to the present are slightly below 1, but they exceed 1 if one compares the Clouds with the Galaxy at times when it had the corresponding metallicities.

Key words: Magellanic Clouds; stars: abundances; stars: mass function; supernovae: general

1 INTRODUCTION

The histories of star formation and chemical evolution in the Magellanic Clouds exhibit distinct features in comparison to those of the disk of the Milky Way. In both Clouds, as reviewed by Olszewski, Suntzeff & Mateo (1996) and Westerglund (1997), it seems that the majority of the stars are under 4 Gyr old, with a sprinkling of older stars. In the LMC there is a sudden rise in the star formation rate (SFR) 2 to 4 Gyr ago (Elson, Gilmore & Santiago 1997; Geisler *et al.* 1997) preceded by either a constant lower SFR (Geha *et al.* 1997) or possibly a virtual gap as manifested by the cluster age distribution (Da Costa 1991; van den Bergh 1991), itself preceded by the formation of a minority older population resembling the Galactic globular clusters although having disk-like kinematics, while in the SMC the SFR appears to have been more uniform (van den Bergh 1991; Olszewski, Suntzeff & Mateo 1996). In the LMC, the sudden rise in the SFR a few Gyr ago is reflected in a corresponding sudden

rise in the metallicity, both from $[\text{Fe}/\text{H}]$ in the clusters (Olszewski *et al.*; Geisler *et al.*) and from α -particle elements in planetary nebulae (Dopita *et al.* 1997), while in the SMC there are fewer data, but some signs of a break in the age-metallicity relation (AMR) at an age between 2 and 4 Gyr (see below). From Olszewski *et al.*, the relative numbers of known clusters older and younger than 3 Gyr are 15:100 in the LMC and 5:37 in the SMC, i.e. about equal proportions, but only one cluster is known in the LMC with age between 3 and 10 Gyr (and that one could be an interloper), while in the SMC there are 3 clusters known in the same interval.

Many models have been put forward for the chemical evolution of the Magellanic Clouds, paying attention to the distinct Fe/O and Fe/ α ratios which are generally found to be higher than in Galactic stars with the same metallicity, i.e. Fe/H. Gilmore & Wyse (1991) pointed out that one way to get this effect is to assume separated star formation bursts, with SNIa contributing extra iron during quiescent intervals. Russell & Dopita (1992) put forward a very de-

tailed and comprehensive pair of models covering several element:element ratios previously determined by themselves and others and normalised to the solar neighbourhood (ISM and F supergiants) rather than the Sun. This model assumed a smooth evolution in which the Clouds are built up by inflow of unprocessed gas according to certain time scales and a slightly steeper IMF than for the solar neighbourhood, which helps to bring down O/Fe and α /Fe. They noted that the observations seem to indicate [O/Fe] slightly lower than [α /Fe], which might be a sign of O/ α stellar yield ratios increasing with mass. They found but minor variations in heavy-element:iron ratios relative to the solar neighbourhood in the LMC, with the exception of an excess of Nd and Sm which have a significant contribution from the r-process. In the SMC they found a still larger excess of Nd and the essentially pure r-process elements Sm and Eu, averaging 0.6 dex, and concluded that the main s-process had been less effective, and the r-process more effective, than in the solar neighbourhood. Russell & Dopita's work referred to young objects (supergiants and the interstellar medium) and therefore did not include discussion of the age-metallicity relation.

Another similarly comprehensive set of models was put forward by Tsujimoto *et al.* (1995), with the express purpose of explaining a relatively high SNIa/SNII ratio, deduced from the Fe/O ratio and supported by some direct counts of remnants, as had been proposed earlier by Barbuy, de Freitas Pacheco & Castro (1994). Tsujimoto *et al.* considered both smooth and bursting models. From an optimal fit to 14 elemental abundances from O up to Ni and smooth (bursting) chemical evolution models they deduced values of 0.15, 0.24 (0.21) and 0.3 (0.28) for the ratio, integrated over time, of SNIa/SN(II+Ib+Ic) in the solar neighbourhood, the LMC and the SMC respectively, and suggested that they originate from Salpeter-like power-law exponents in the IMF of respectively 1.33, 1.71 (1.62) and 1.88 (1.84). These steeper IMFs (between fixed upper and lower mass limits), also postulated by Russell & Dopita, then also serve to produce the subsolar metallicities in the Magellanic Clouds at the corresponding gas fractions and give reasonable age-metallicity relations, especially with the burst models.

The hypothesis of a slightly steeper IMF compared to the Galaxy is neither supported nor ruled out by direct star counts (Hill, Madore & Freedman 1994 and references therein). An alternative way to explain low metallicities is to assume outflow (e.g. Carigi *et al.* 1995). Such outflow can be assumed to be either homogeneous or selective, the latter case being associated with starbursts and leading to enhancement of the Fe/O ratio, among others (e.g. Marconi, Matteucci & Tosi 1994). Such a model has been put forward for the LMC by Pilyugin (1996), who points out that a closed bursting model, like those of Gilmore & Wyse, will produce the same final oxygen abundance at a given gas fraction as a smooth model, so that some other factor is needed to provide the reduction in metallicity compared to the solar neighbourhood with the same gas fraction. Given the absence of compelling evidence for a change in the IMF, Pilyugin postulates selective winds associated with star formation bursts; an initial square-wave burst in the first 0.2 Gyr, accounting for 8 per cent of the stars, is followed by a hiatus until 9 Gyr, when a major burst or series of bursts begins, ending after 12.5 Gyr (0.5 Gyr before present) and

accounting for the remainder, in accordance with the age distribution of the LMC clusters. A model with both homogeneous and selective galactic winds (with a selective expulsion of 75 per cent of ejecta from massive stars and a homogeneous expulsion of a mass equal to that being made into stars) is found to give good fits to the AMR and to [Fe/O].

Unfortunately, the assessment of [Fe/O] and, to a lesser extent, other abundance ratios is dependent on which Galactic standards are used (cf Pagel 1992). While it is certainly true that [Fe/O] is not negative in the Clouds, it is not strongly positive either when compared to Galactic supergiants instead of the Sun, and because of various systematic effects the supergiants may provide a better standard. Among α -particle elements, notably Mg and Ca, and for Ti which seems to behave like the α -elements among Galactic stars, the offset from either solar or local supergiant ratios is insignificantly different from zero, compared to small positive values in the Milky Way for the same [Fe/H] (Edvardsson *et al.* 1993). There is thus some difference, but not necessarily large enough to demand either a changed IMF or selective galactic winds. Some sort of winds are probably present, as witness the Magellanic stream, and these may well account for the relatively low metallicities, and we agree with Russell & Dopita and with Tsujimoto *et al.* in assuming a significant effect of inflow, since this is needed, at least in smooth models with reasonable star formation laws, to account for the 'G-dwarf' problem posed by the relatively small number of old objects. The existence of separated star formation bursts is well attested in the LMC by population studies of both clusters and field stars, and is supported in both Clouds by the AMRs.

With this background, it seems worth while to investigate how well the chemical evolution of the Magellanic Clouds can be modelled with yields and time-delays equal to those applying to the solar neighbourhood, but with inflow and a homogeneous wind. In two previous papers (Pagel & Tautvaišienė 1995, 1997) we developed analytical models giving a fair fit to stellar abundances of primary elements in the solar neighbourhood, and in this paper we shall attempt to apply the same ideas to the Clouds.

2 THE MODEL

2.1 General description

In agreement with Russell & Dopita (1992) and Tsujimoto *et al.* (1995), we assume the Clouds to have been built up by gradual infall of unprocessed material. This helps to alleviate the 'G-dwarf' problem that would otherwise arise from the small relative proportion of old, metal-poor stars. Our formalism is an adaptation of that of Pagel (1997) as detailed below; like Tsujimoto *et al.*, we assume linear laws of star formation and investigate both smooth and bursting models. Like Pilyugin (1996), we assume yields and time delays identical to those which apply to the solar neighbourhood and appeal to galactic winds to explain the low metallicities of the Clouds in relation to their current gas fractions; we take the specific numbers from our two previous papers. Unlike Pilyugin, however, we ignore selective winds and assume just a non-selective wind proportional to the SFR, similar to the model of Hartwick (1976) for the Galactic halo.




Figure 1. SFR history for the LMC according to our model. The full curve shows the bursting model according to Table 1, while the broken-line curve shows a smooth model with $u = 0.18t$.




Figure 2. SFR history for the SMC according to our model. The full curve shows the bursting model according to Table 2, while the broken-line curve shows a smooth model with $u = 0.115t$.

2.2 The formalism

We assume a linear or quasi-linear star formation law

$$\frac{ds}{du} = g, \quad (1)$$

where s is the mass in stars (+ remnants), g is the mass of gas and $u = \int_0^t \omega(t') dt'$ where t is time and ω is the inverse time scale for star formation, assumed constant in the ‘smooth’ models. In ‘bursting’ models, ω is assumed to be constant over certain time periods, between which it changes discontinuously. Inflow is assumed to occur at a rate

$$f(t) = \omega(t) e^{-u}, \quad (2)$$




Figure 3. Age-abundance relation for α -elements in the LMC. Curves are as in Fig. 1. Data points are from Dopita *et al.* (1997): open circles represent oxygen while filled circles represent an average of Ne, S and Ar.

and outflow at a rate

$$e(t) = \eta \frac{ds}{dt} = \eta \omega g, \quad (3)$$

with $\eta = \text{const.}$, so that the gas mass and total mass satisfy the differential equations

$$\frac{dg}{du} = e^{-u} - (1 + \eta)g \quad (4)$$

and

$$\frac{dm}{du} = e^{-u} - \eta g \quad (5)$$

respectively. From Eqs. (4), (5), the gas and total masses evolve according to

$$g(u) = [e^{-u} - e^{-(1+\eta)u}] / \eta; \quad (6)$$

$$m(u) = [1 - e^{-(1+\eta)u}] / (1 + \eta); \quad (7)$$

the star mass is

$$s(u) = m(u) - g(u) \quad (8)$$

and the gas fraction is

$$\mu(u) = g(u)/m(u). \quad (9)$$

Using the instantaneous recycling approximation, the abundance of a promptly produced primary element such as O or Mg, in units of its yield (assumed constant), satisfies

$$\frac{d}{du}(gz_1) = g[1 - (1 + \eta)z_1], \quad (10)$$

which has the solution

$$z_1(u) = \frac{1}{\eta} - \frac{u}{e^{\eta u} - 1}. \quad (11)$$

Results of Eqs. (1) to (11) for our assumed model parameters are given in the first seven columns of Tables 1 and 2. Figs 1 and 2 show the resulting SFR histories for both smooth and bursting models; the rough coincidence in time between our assumed bursts in the two Clouds suggests that they may have resulted from some mutual interaction. For the LMC, our assumed value of 1 for η agrees with that adopted by Pilyugin and the current star formation rate is about average; this may be compared with the remark by Westerlund (1997) that the current SFRs in both Clouds deduced from H α emission may indicate that their current SFRs are below their averages. Thus the decline that we assume in the last 3 Gyrs from the peak of the recent burst, resulting from our assumption of Eq. (1) with a constant ω in the relevant interval, seems to be fairly realistic. The final gas fraction adopted for the SMC is in accordance with the number given by Westerlund; for the LMC it is somewhat larger than his figure of $< 8\%$, but in accordance with that adopted in other models in the literature and similar to some estimates for the solar neighbourhood. Fig 3 shows the age-abundance relation in the LMC for O, Ne, S and Ar, assumed to be instantaneously produced with a yield of 0.7 times their solar abundance, and confirms that a bursting model gives a better fit than a smooth one.

For elements (or components thereof) that are ejected after a significant time delay, we use the ‘delayed production’ approximation (Pagel 1989) which assumes the element to be ejected at a fixed time delay Δ after the time of star formation, whereas the total mass ejection from a generation

Table 1. Bursting model for the LMC; $\langle\omega\rangle = 0.18 \text{ Gyr}^{-1}$; $\eta = 1$.

ω Gyr^{-1}	t Gyr	u	g	m	μ	z_1	z_2 0.023	z_2 0.037	z_2 1.33	z_2 2.67	[O/H]	[Fe/H]
0.15	0.023	0.003	0.003	0.003	1.00	0.002	0.000	0.000	0.000	0.000	-2.92	-3.31
	0.037	0.006	0.006	0.006	1.00	0.003	0.000	0.000	0.000	0.000	-2.71	-3.11
	0.500	0.075	0.067	0.070	0.96	0.037	0.034	0.032	0.000	0.000	-1.59	-1.98
	0.900	0.135	0.110	0.118	0.93	0.066	0.063	0.061	0.000	0.000	-1.34	-1.73
	1.330	0.200	0.148	0.165	0.90	0.096	0.094	0.092	0.000	0.000	-1.17	-1.57
0.15	2.000	0.300	0.192	0.226	0.85	0.143	0.140	0.139	0.022	0.000	-1.00	-1.31
0.08	2.023	0.302	0.193	0.227	0.85	0.143	0.143	0.141	0.024	0.000	-1.00	-1.30
	2.037	0.303	0.193	0.227	0.85	0.144	0.143	0.143	0.025	0.000	-1.00	-1.30
	2.670	0.354	0.209	0.254	0.83	0.166	0.166	0.165	0.073	0.000	-0.93	-1.11
	3.330	0.406	0.222	0.278	0.80	0.190	0.189	0.188	0.135	0.007	-0.88	-0.96
	4.670	0.514	0.240	0.321	0.75	0.235	0.234	0.234	0.184	0.046	-0.78	-0.84
	6.000	0.620	0.249	0.355	0.70	0.278	0.278	0.277	0.234	0.116	-0.71	-0.75
	7.000	0.700	0.250	0.377	0.66	0.310	0.309	0.309	0.270	0.167	-0.66	-0.70
	8.000	0.780	0.248	0.395	0.63	0.340	0.339	0.339	0.306	0.215	-0.62	-0.65
	9.000	0.860	0.244	0.411	0.59	0.369	0.369	0.369	0.341	0.261	-0.59	-0.61
	10.000	0.940	0.238	0.424	0.56	0.397	0.397	0.397	0.375	0.306	-0.56	-0.57
0.08	11.000	1.020	0.231	0.435	0.53	0.425	0.425	0.425	0.408	0.349	-0.53	-0.54
0.50	11.023	1.032	0.229	0.437	0.53	0.429	0.419	0.419	0.402	0.345	-0.52	-0.54
	11.037	1.039	0.229	0.437	0.52	0.431	0.421	0.415	0.399	0.342	-0.52	-0.54
	11.500	1.270	0.202	0.461	0.44	0.504	0.498	0.495	0.320	0.280	-0.45	-0.56
	12.330	1.685	0.151	0.483	0.31	0.616	0.616	0.615	0.256	0.236	-0.37	-0.55
	12.500	1.770	0.141	0.486	0.29	0.637	0.637	0.637	0.359	0.234	-0.35	-0.48
	12.670	1.855	0.132	0.488	0.27	0.007	0.657	0.657	0.455	0.234	-0.35	-0.43
	13.000	2.020	0.115	0.491	0.23	0.691	0.693	0.695	0.626	0.239	-0.32	-0.34
	13.670	2.355	0.086	0.496	0.17	0.753	0.757	0.760	0.915	0.270	-0.28	-0.23
	13.800	2.420	0.081	0.496	0.16	0.764	0.769	0.771	0.963	0.422	-0.27	-0.21
0.50	14.000	2.520	0.074	0.497	0.15	0.780	0.785	0.787	1.033	0.645	-0.26	-0.19

Table 2. Bursting model for the SMC; $\langle\omega\rangle = 0.115 \text{ Gyr}^{-1}$; $\eta = 2$.

ω Gyr^{-1}	t Gyr	u	g	m	μ	z_1	z_2 0.023	z_2 0.037	z_2 1.33	z_2 2.67	[O/H]	[Fe/H]
0.10	0.023	0.002	0.002	0.002	1.00	0.001	0.000	0.000	0.000	0.000	-3.10	-3.49
	0.037	0.004	0.004	0.004	1.00	0.002	0.000	0.000	0.000	0.000	-2.89	-3.29
	0.500	0.050	0.045	0.046	0.97	0.025	0.023	0.021	0.000	0.000	-1.76	-2.16
	1.000	0.100	0.082	0.086	0.95	0.048	0.046	0.045	0.000	0.000	-1.47	-1.87
0.10	1.330	0.133	0.102	0.110	0.93	0.064	0.062	0.061	0.000	0.000	-1.35	-1.75
0.01	1.353	0.133	0.102	0.110	0.93	0.064	0.064	0.063	0.000	0.000	-1.35	-1.75
	1.367	0.133	0.102	0.110	0.93	0.064	0.064	0.064	0.000	0.000	-1.35	-1.75
	2.660	0.146	0.110	0.118	0.93	0.070	0.070	0.070	0.067	0.000	-1.31	-1.32
	4.000	0.160	0.117	0.127	0.92	0.076	0.076	0.076	0.072	0.002	-1.28	-1.29
	6.000	0.180	0.126	0.139	0.91	0.085	0.085	0.085	0.081	0.019	-1.23	-1.24
	8.100	0.201	0.135	0.151	0.90	0.094	0.094	0.094	0.089	0.034	-1.18	-1.20
0.01	10.000	0.220	0.143	0.161	0.89	0.102	0.102	0.102	0.097	0.047	-1.15	-1.16
0.35	10.023	0.228	0.146	0.165	0.88	0.105	0.098	0.098	0.093	0.045	-1.13	-1.16
	10.037	0.233	0.147	0.168	0.88	0.107	0.100	0.095	0.091	0.044	-1.12	-1.17
	10.200	0.290	0.165	0.194	0.85	0.131	0.124	0.121	0.070	0.035	-1.04	-1.18
	10.500	0.395	0.184	0.231	0.80	0.172	0.167	0.164	0.048	0.024	-0.92	-1.17
	11.000	0.570	0.192	0.273	0.70	0.232	0.229	0.227	0.030	0.017	-0.79	-1.11
	11.330	0.685	0.188	0.291	0.65	0.267	0.265	0.263	0.024	0.014	-0.73	-1.07
	12.000	0.920	0.168	0.312	0.54	0.326	0.326	0.326	0.189	0.012	-0.64	-0.77
	12.670	1.154	0.142	0.323	0.44	0.373	0.373	0.374	0.337	0.012	-0.58	-0.61
	13.300	1.375	0.118	0.328	0.36	0.406	0.408	0.409	0.451	0.244	-0.55	-0.52
0.35	14.000	1.620	0.095	0.331	0.29	0.434	0.436	0.438	0.550	0.494	-0.52	-0.45




Figure 4. Age-metallicity relation for the LMC. Curves represent our models as in previous figures. Data sources are as follows: *open circles*, Olszewski *et al.* (1991); *crosses*, Girardi *et al.* (1995); *open squares*, Geisler *et al.* (1997); *filled circles*, Idiart & de Freitas Pacheco (1997).




Figure 5. Age-metallicity relation for the SMC. Curves represent our models as in previous figures. Data represented by open circles are from Da Costa (1991).

of stars is still assumed to take place instantaneously. In this case the abundance z_2 in units of the yield varies according to

$$\frac{d}{dt}(gz_2) = \omega'g' - (1 + \eta)z_2\omega g; \quad t > \Delta, \quad (12)$$

where the dashes indicate that the variable has to be taken at the time $t - \Delta$ rather than t . Eq. (12) leads after some reduction to

$$\frac{d}{dt}[z_2(e^{\eta u} - 1)] = \omega'e^{(1+\eta)u}[e^{-u'} - e^{-(1+\eta)u'}]. \quad (13)$$

For smooth models ($\omega = \text{const.}$), Eq. (13) has the solution

$$z_2(u) = \frac{e^{(1+\eta)\omega\Delta}}{e^{\eta u} - 1} \left[\frac{e^{\eta(u-\omega\Delta)} - 1}{\eta} - (u - \omega\Delta) \right]; \quad u > \omega\Delta. \quad (14)$$

In bursting models, we assume $\omega = \omega_1$, say, up to time t_1 , $\omega = \omega_2$ between t_1 and t_2 and $\omega = \omega_3$ between t_2 and $t_3 = T$, the age of the system, assumed to be 14 Gyr. ω_2 is small, representing a quiescent phase between an initial starburst represented by ω_1 and a stronger recent starburst represented by ω_3 . Then, in Eq. (13),

$$\omega' = u' = 0; \quad \text{for } t < \Delta; \quad (15)$$

$$\begin{aligned} \omega' &= \omega_1; \\ u' &= \omega_1(t - \Delta); \quad \text{for} \end{aligned}$$

$$\Delta \leq t < t_1 + \Delta; \quad (16)$$

$$\begin{aligned} \omega' &= \omega_2; \\ u' &= \omega_1 t_1 + \omega_2(t - t_1 - \Delta); \quad \text{for} \\ t_1 + \Delta &\leq t < t_2 + \Delta; \end{aligned} \quad (17)$$

$$\begin{aligned} \omega' &= \omega_3; \\ u' &= \omega_1 t_1 + \omega_2(t_2 - t_1) + \omega_3(t - t_2 - \Delta); \quad \text{for} \\ t &\geq t_2 + \Delta. \end{aligned} \quad (18)$$

Also

$$\begin{aligned} u &= \omega_1 t; \quad \text{for } t < t_1; \\ &= \omega_1 t_1 + \omega_2(t - t_1); \quad \text{for } t_1 \leq t < t_2; \\ &= \omega_1 t_1 + \omega_2(t_2 - t_1) + \omega_3(t - t_2); \quad \text{for } t \geq t_2. \end{aligned} \quad (19)$$

Thus in separate time segments Δ to t_1 (or t_1 to Δ if $\Delta > t_1$), t_1 (or Δ) to $t_1 + \Delta$, $t_1 + \Delta$ to t_2 , t_2 to $t_2 + \Delta$, $t_2 + \Delta$ to $t_3 = T$, $\omega' = \text{const.}$ and the exponents in Eq. (13) are all linear functions of time:

$$\begin{aligned} (1 + \eta)u &= p + qt; \\ u' &= r + st, \end{aligned} \quad (20)$$

say, and so Eq. (13) can be solved analytically for each interval t_a to t_b :

$$\begin{aligned} &z_2(t_b)(e^{\eta u_b} - 1) - z_2(t_a)(e^{\eta u_a} - 1) \\ &= \omega' \frac{e^{p-r}}{q-s} \{ e^{(q-s)t_b} - e^{(q-s)t_a} \} \\ &- \omega' \frac{e^{p-(1+\eta)r}}{q-(1+\eta)s} \{ e^{[q-(1+\eta)s]t_b} - e^{[q-(1+\eta)s]t_a} \}. \end{aligned} \quad (21)$$

2.3 Yields and time delays



Table 3 gives the yields and time delays that we assume, after Pagel & Tautvaišienė (1995, 1997). Yields are in units of solar abundance of the corresponding element and time delays in Gyr are given in the top row. Total abundances are obtained for each time step by summing the products of these yields with the appropriate values of z for the relevant time delay, which are given in columns 7 to 11 of Tables 1 and 2. Figs 4 and 5 show the resulting age-metallicity relations for the two Clouds, which are similar to those derived by Tsujimoto *et al.* (1995), but differ somewhat from those for the LMC by Pilyugin (1996).

3 ELEMENT:ELEMENT RATIOS

The last seven columns of Tables 1 and 2 give abundances or constituent z -values for individual elements in the Clouds calculated from our bursting models, and element:iron ratios are plotted in Figs 6 to 11. The observational data, which come mainly from MC supergiants, have in some cases been adjusted to be relative to Galactic supergiants rather than the Sun, since (as noted by Russell & Dopita 1992, Pagel 1992 and Hill, Andrievsky & Spite 1995) there are some discrepancies, particularly in the case of oxygen, and it seems better to compare like with like. Russell & Dopita took $[\text{O}/\text{Fe}]$ for the solar neighbourhood to be -0.03 only, but various studies of Galactic supergiants (Luck & Lambert

Table 3. Yields and time delays

Δ (Gyr)	0	0.023	0.037	1.33	2.67
O	0.70				
Mg	0.88				
Si, Ti	0.70			0.12	
Ca	0.56			0.18	
Fe	0.28			0.42	
Sr, La	0.013	0.11	0.29		0.29
Y, Ba	0.010	0.08	0.30		0.30
Zr	0.022	0.17	0.28		0.28
Ce	0.020	0.16	0.26		0.26
Pr, Sm	0.055	0.46	0.11		0.11
Nd	0.040	0.33	0.17		0.17
Eu, Dy	0.080	0.64	0.01		0.01

**Figure 6.** Element:iron ratios for oxygen, α -particle elements and Ti in the LMC and in the anomalous Galactic halo stars. The full-drawn curves show predictions from the bursting model and the broken-line curves those from the smooth model. Data points for the LMC stars have been shifted upwards by 0.2 dex for oxygen and downwards by 0.1 dex for silicon. Data sources: + *signs*, Luck & Lambert (1992); *open circles*, Thévenin (1997); *five-cornered stars*, Hill, Andriewski & Spite (1995); *open squares*, Spite, Barbuy & Spite (1993); *open diamonds*, Jüttner et al. (1992); *open triangles*, McWilliam & Williams (1991); *small six-pointed stars*, Barbuy, de Freitas Pacheco & Castro (1994); *asterisks*, Richtler, Spite & Spite (1989); *crosses*, Russell & Bessell (1989); *large six-pointed stars*, anomalous Galactic halo stars after Nissen & Schuster (1997).**Figure 7.** Element:iron ratios for s-process elements in the LMC and in the anomalous Galactic halo stars. Curves and data sources as in Fig 6. Data points for Ba and La in the LMC stars have been shifted downwards by 0.1 dex and 0.16 dex respectively.

1985; Spite, Barbuy & Spite 1989; Hill, Andrievsky & Spite 1995; Venn 1995) indicate a more negative value, $[O/Fe] \approx -0.2$ relative to the Sun. The observational $[O/Fe]$ ratios plotted in Figs 6 and 9 have accordingly been shifted upwards by 0.2 dex, which leads to a somewhat different picture from that assumed in previous chemical evolution models for the Clouds; in particular, there is now (apparently) good agreement between oxygen and the other α -elements. The oxygen abundances from planetary nebulae shown in Fig 2 also support this upward adjustment. On the basis of studies of Canopus and other Galactic supergiants by Luck (1982), Reynolds, Hearnshaw & Cottrell (1988), Russell & Bessell (1989) and Spite, Spite & François (1989), we have shifted the data points for $[Si/Fe]$, $[Ba/Fe]$ and $[Nd/Fe]$ downwards by 0.1 dex and $[La/Fe]$ downwards by 0.16 dex, taking all the remaining elements at face value.

Although the model curves in Figs 6 to 11 are meant to represent an evolution in time, it has to be borne in mind that the data points come from young supergiants, so that their horizontal spread is due to scatter (part of which is probably observational) and not from evolution in time. Consequently it is only the centroid of these points that is significant and no evolutionary trend can be deduced from the data. However, Nissen & Schuster (1997) have found a sample of stars from the outer halo of our Galaxy in which, among other peculiarities, the α -element to iron ratios are




Figure 8. Element:iron ratios for heavy metals in the LMC. Curves and data sources as in Fig 6. Data points for Nd have been shifted downwards by 0.1 dex.

more or less solar rather than enhanced as in other stars with the same metal deficiency. They suggest that these stars could have been captured from dwarf galaxies such as the Magellanic Clouds and we accordingly include them (without any adjustments) in the relevant plots. Our LMC model fits these ratios very well, including a slight rise from solar-like $[\alpha/\text{Fe}]$ around $[\text{Fe}/\text{H}] = -0.8$ to $[\alpha/\text{Fe}] \simeq 0.2$ at $[\text{Fe}/\text{H}] = -1.2$ (Fig 6) and a more uniform trend for s-process elements (Fig 7), supporting an origin in the LMC (or some similar system). The fit of our SMC model (Figs 9, 10) is not quite as good. As far as the actual Magellanic stars are concerned, there is considerable scatter in the data, much of which we suspect to be unreal, although we have chosen to plot individual determinations rather than to follow Russell & Dopita in just plotting an average. Within the scatter, the agreement of our model with abundances deduced from observation seems to be quite satisfactory, with the possible exception of a few of the heaviest elements. Broadly speaking, there is hardly any evidence for significant departures from solar, or solar-neighbourhood, abundance ratios among the elements considered (we do not discuss carbon, nitrogen or sodium which all exhibit ‘secondary’ behaviour), and that result is just what is predicted; whereas the steepened IMF models of Tsujimoto *et al.*, while successful for oxygen (if unadjusted to Galactic supergiants), predict sub-solar abundances (relative to iron) of several α -particle elements, notably magnesium, which are not supported by




Figure 9. Element:iron ratios for oxygen, α -particle elements and titanium for the SMC and anomalous Galactic halo stars. Data points for oxygen and silicon in the SMC have been shifted upwards by respectively 0.2 dex and 0.1 dex. Solid and broken curves represent our SMC bursting and smooth models respectively. Data sources: *open diamonds*, Jüttner *et al.* (1992); *filled diamonds*, Hill, Barbuy & Spite (1997); *squares with + signs*, Spite, Barbuy & Spite (1989); *open squares*, Hill & Spite (1997); *open triangles*, Spite *et al.* (1986); *open circles*, Spite, Richtler & Spite (1991); *filled circles*, Thévenin (1997); *+ signs*, Luck & Lambert (1992); *crosses*, Russell & Bessell (1989); *asterisks*, Spite, Spite & François (1989); *filled triangles*, Hill (1997); *filled five-pointed stars*, Thévenin & Foy (1986); *open five-pointed stars*, Foy (1981); *open six-pointed stars*, anomalous Galactic halo stars from Nissen & Schuster (1997).

the data. Pending improved data, especially for the heaviest elements, we conclude that neither a steepened IMF nor selective galactic winds are required to explain the abundances in the Magellanic Clouds.

4 RELATIVE SUPERNOVA RATES

Barbuy, de Freitas Pacheco & Castro (1994) and Tsujimoto *et al.* (1995) have claimed that the overall ratios of Type Ia supernovae to core-collapse supernovae ever formed in the Clouds are higher than in the solar neighbourhood. Our models do not support this and they fall down if it is true. With our approximations, the ratio is proportional to

$$\frac{s(T - \Delta)}{s(T)} = \frac{m(T - \Delta) - g(T - \Delta)}{m(T) - g(T)}. \quad (22)$$




Figure 10. Element:iron ratios for s-process elements in the SMC and anomalous Galactic halo stars. Curves and symbols as in Fig. 9. Data points for Ba and La in the SMC stars have been shifted downwards by 0.1 dex and 0.16 dex respectively.




Figure 11. Element:iron ratios for heavy metals in the SMC. Curves and symbols as in Fig. 9. Data points for Nd have been shifted downwards by 0.1 dex.

Taking $u = 4.5$, $\omega\Delta = 0.4$ for the solar neighbourhood (Pagel & Tautvaišienė 1995), this ratio comes out to be 0.95, whereas in the LMC and SMC according to Tables 1 and 2 it is 0.86 and 0.77 respectively, i.e. marginally lower. The situation changes if we compare the above Magellanic ratios with those that prevailed in the solar neighbourhood at those times in the past when it had the corresponding metallicities; these turn out to be 0.81 for $[\text{Fe}/\text{H}] = -0.2$ and 0.62 for $[\text{Fe}/\text{H}] = -0.45$. Thus we attribute the lower α/Fe ratios chiefly to the effect of slower evolution up to a metallicity which prevailed in our Galaxy at an earlier time when there had indeed been relatively fewer SNIa in the latter. It can be seen from Figs 6 to 11 that the influence of starbursts on the element:element ratios is comparatively insignificant in our models.

When we consider *current* supernova rates, the situation is quite different again. In this case the ratio is proportional to

$$\frac{(ds/dt)_{T-\Delta}}{(ds/dt)_T} = \frac{g(T-\Delta)}{g(T)}. \quad (23)$$

The corresponding numbers are 1.2 for the solar neighbourhood, 1.8 for the LMC and 1.5 for the SMC in our bursting models. Thus we expect a 50 per cent higher ratio of SNIa to core-collapse supernovae in the LMC at present. Assuming a ratio in the solar neighbourhood between 0.11 and 0.25 (van den Bergh & McClure 1994), the ratio in the LMC is then predicted to be between 0.16 and 0.38, which is in good agreement with the average figure of 0.29 given for Sdm-Im galaxies by Cappellaro *et al.* (1993). A still higher ratio, of order 1, has been estimated for the LMC on the basis of X-ray observations of remnants by Hughes *et al.* (1995), who suggest a likely lower limit of 0.25, so that our model agrees qualitatively, if not quantitatively, with their findings. Current ratios are sensitive to the precise assumptions made about the bursts, and do not necessarily represent historic averages. We suggest that the relatively high ratio of SNIa to core collapse SN observed in the LMC and other Sd-Im galaxies is related to their star formation history rather than the IMF.

ACKNOWLEDGMENTS

GT acknowledges a NORDITA Baltic/NW Russia Fellowship from the Nordic Council of Ministers and hospitality of NORDITA while the work for this paper was being carried out.

REFERENCES

- Barbuy B., de Freitas Pacheco J. A., Castro S. 1994, A&A, 283, 32
- Cappellaro E., Turatto M., Benetti S., Tsvetkov D.Yu., Bartunov O.S., Makarova I.N. 1993, A&A, 273, 383
- Carigi L., Colín P., Peimbert M. & Sarmiento A. 1995, ApJ, 445, 98
- Da Costa G. S. 1991, in The Magellanic Clouds, eds. Haynes R., Milne D., IAU Symp. 148, Kluwer, p. 183
- Dopita M. A., Vassiliadis E., Wood P. R., Meatheringham S. J., Harrington J. P., Bohlin R. C., Ford H. C., Stecher T. P., Maran S. P. 1997, ApJ, 474, 188

- Edvardsson B., Andersen J., Gustafsson B., Lambert D.L., Nissen P.E., Tomkin J. 1993, A&A, 275, 101
- Elson R.W., Gilmore G.F., Santiago B.X. 1997, MNRAS, 289, 157
- Foy R., 1981, A&A, 103, 135
- Geha M.C., Holtzman J.A., Mould J.R., Gallger J.S. III *et al.* 1997, astro-ph 9711144
- Geisler D., Bica E., Dottori H., Claría J.J., Piatti A.E., Santos J.F.C., Jr. 1997, AJ, 114, 1920
- Gilmore G., Wyse R.F.G. 1991, ApJ, 367, L55
- Girardi L., Chiosi C., Bertelli G., Bressan A. 1995, A&A, 298, 87
- Hartwick F.D.A. 1976, ApJ, 209, 418
- Hill R.J., Madore B.F., Freedman W.L. 1994, ApJ, 429, 192
- Hill V. 1997, A&A, 324, 435
- Hill V., Andrievsky S., Spite M. 1995, A&A, 293, 347
- Hill V., Barbuy B., Spite M. 1997, A&A, 323, 461
- Hill V., Spite M. 1997, Poster at IAU Assembly, Kyoto
- Hughes J.P., Hayashi I., Helfand D., Hwang U., Itoh M., Kirshner R., Koyama K., Markert T., Tsunemi H., Woo J. 1995, ApJ, 444, L81.
- Idiart T. P., de Freitas Pacheco J. A. 1997, The Initial Mass Function, Herstmonceux Conference, ed. G. Gilmore
- Jüttner A., Stahl O., Wolf B., Baschek B. 1992, in New Aspects of Magellanic Cloud Research, eds. Baschek B., Klare G., Lequeux J., Springer-Verlag, p. 337
- Luck R. E. 1982, ApJ, 263, 215
- Luck R. E., Lambert D. L. 1985, ApJ, 298, 782
- Luck R. E., Lambert D. L. 1992, ApJS, 79, 303
- McWilliam A., Williams R. E. 1991, in The Magellanic Clouds, IAU Symp. 148, eds. Hayes R., Milne D., Kluwer, p. 391
- Marconi G., Matteucci F., Tosi M. 1994, MNRAS, 270, 35
- Nissen P.E., Schuster W.J. 1997, A&A, 326, 751
- Olszewski E.W., Suntzeff N.B., Mateo M. 1996, ARAA, 34, 511
- Olszewski E. W., Schommer R. A., Suntzeff N. B., Harris, H. C. 1991, AJ, 101, 515
- Pagel B.E.J. 1989, Rev. Mex. Astr. Astrofis., 18, 153
- Pagel B.E.J. 1992, New Aspects of Magellanic Cloud Research, eds. B. Baschek, J. Klare, J. Lequeux, Springer-Verlag, p. 330
- Pagel B.E.J. 1997, *Nucleosynthesis and Chemical Evolution of Galaxies*, Cambridge University Press, Chapter 8
- Pagel B.E. J., Tautvaisienė G. 1995, MNRAS, 276, 505
- Pagel B.E.J., Tautvaisienė G. 1997, MNRAS, 288, 108
- Pilyugin L. S. 1996, A&A, 310, 751
- Reynolds S. E., Hearnshaw J. B., Cottrell P. L. 1988, MNRAS, 235, 1423
- Richtler T., Spite M., Spite F. 1989, A&A, 225, 351
- Russell S. C., Bessell M. S. 1989, ApJS, 70, 865
- Russell S.C., Dopita M.A. 1992, ApJ, 384, 508
- Spite M., Barbuy B., Spite F. 1989, A&A, 222, 35
- Spite M., Barbuy B., Spite F. 1993, A&A, 272, 116
- Spite M., Cayrel R., François P., Richtler T., Spite F. 1986, A&A, 168, 197
- Spite F., Spite M., François P. 1989, A&A, 210, 25
- Spite F., Richtler T., Spite M. 1991, A&A, 252, 557
- Thévenin F. 1997, CDS Bull., submitted
- Thévenin F., Foy R. 1986, A&A, 155, 145
- Tsujimoto T., Nomoto K., Yoshii Y., Hashimoto M., Yanagida S., Thielemann F.-K. 1995, MNRAS, 277, 945
- van den Bergh S. 1991, ApJ, 369, 1
- van den Bergh S., McClure, R.D. 1994, ApJ, 425, 205
- Venn K. A., ApJ, 1995, 449, 839
- Westerlund B. 1997, *The Magellanic Clouds*, Cambridge University Press

This paper has been produced using the Royal Astronomical Society/Blackwell Science L^AT_EX style file.